

## **WHAT TO EXPECT AFTER SCAPHOID SURGERY**

**Dr. Erik Dorf**

Scaphoid fractures can be complex injuries, however, this type of surgery is common to my practice and I have found the results to be consistent and reliable. Keep in mind that every person responds differently to surgery and treatment, and your recovery may vary from the below outline. Typical factors which influence the recovery process include:

- Your age and health.
- The severity of your fracture/nature of your injury
- The length of time between the injury and the surgery
- Your compliance with prescribed treatment and therapy



1. ***SURGERY DAY:*** I will give you a copy of the x-ray of your surgically repaired hand. You will be placed in a short splint that immobilizes your thumb, covered by an ace wrap. Your remaining fingers and elbow will move freely. You will keep this on until your first follow up visit, which occurs at two weeks after surgery. Please do not remove the splint or get it wet. You may use a kitchen garbage bag and duct tape to keep it dry in the shower/bathtub.
2. ***FIRST 48 HOURS AFTER SURGERY:*** You should try **to elevate your hand and wiggle your fingers** as much as possible. Ice can also be used to reduce inflammation. You will be provided with pain medication. I discourage the use of ibuprofen for the first two weeks after surgery.
3. ***FIRST TWO WEEKS:*** After surgery you should wiggle your fingers (not your thumb) as much as possible. **No heavy lifting.**
4. ***AT YOUR TWO WEEK POST OP OFFICE VISIT:*** We will place you into a cast. We do not get x-rays at your first visit. You are again encouraged to use your hand as much as possible after being placed in a cast. Again **no heavy lifting.**
5. ***SIX WEEKS AFTER SURGERY:*** We will remove your cast. We will get x-rays at this time to monitor healing. We will either place you back in a cast or transition you to a removable brace. If you go into a brace, you will receive instructions about how much to wear it and when to wean out of it.
6. ***TEN TO TWELVE WEEKS AFTER SURGERY:*** We will see you back in our clinic for a repeat exam. We will repeat x-rays at that time as well.
7. ***THERAPY:*** If at any time during this treatment course it appears you will require therapy to help decrease swelling, or improve range of motion you will be provided with a prescription for therapy. Approximately half of our patients require some sort of therapy.
8. ***HOW LONG TO GET BETTER?*** **Most** people are **mostly** better by 12 weeks after surgery. No one is completely better at this time, but you can typically return to most of your normal activities.
9. ***HOW LONG FOR FULL RECOVERY?*** It usually takes people **six months to a year** before they can expect full improvement after having surgery.

**Should you have any questions, please call any of our offices at the following numbers:**  
Edwards (970) 569-3240      Frisco (970) 668-3633      Vail (970) 476-7220